

Publication List

- 1 Kambara T, Yasaka T, and Nakamura T.
The role of polymorphonuclear leukocytes in delayed hypersensitivity skin reactions: suppressive effects of anti-polymorphonuclear leukocyte serum.
Virchows Arch. [Cell. Pathol.] 1981; 37: 191-198.
- 2 Kambara T, Kutsuna T, Okamoto T, and Nakamura T.
Inflammatory activities of synovial fluids from rheumatoid- and osteo-arthritis to guinea pig skin.
Acta Pathol. Jpn. 1982; 32: 575-583.
- 3 Kambara T, Kawagoe T, and Nakamura T.
An alkaline proteinase which has inflammatory activity isolated from guinea pig lymphoid cells.
Biochim. Biophys. Acta 1982; 716: 224-231.
- 4 Nakamura T, Yamamoto T, Ishimatsu T, and Kambara T.
A high-molecular-weight trypsinlike protease in the skin sites of delayed hypersensitivity in guinea pigs.
Am. J. Pathol. 1984; 114: 250-263.
- 5 Nakamura T, Board PG, Matsushita K, Tanaka H, Matsuyama T, and Matsuda T.
 α 1-Acid glycoprotein expression in human leukocytes: possible correlation between α 1- acid glycoprotein and inflammatory cytokines in rheumatoid arthritis.
Inflammation 1993; 17: 33-45.
- 6 Nakamura T, Suchard SJ, Abe A, Shayman JA, and Boxer LA.
Role of diradylglycerol formation in H₂O₂ and lactoferrin release in adherent human polymorphonuclear leukocytes.
J. Leukocyte Biol. 1994; 56: 105-109.
- 7 Suchard SJ, Nakamura T, Abe A, Shayman JA, and Boxer LA.
Phospholipase D-mediated diradylglycerol formation coincides with H₂O₂ and lactoferrin release in adherent human neutrophils.
J. Biol. Chem. 1994; 269: 8063-8068.
- 8 Nakamura T, Abe A, Balazovich KJ, Wu D, Suchard JD, Boxer LA, and Shayman JA.
Ceramide regulates oxidant release in adherent human neutrophils.
J. Biol. Chem. 1994; 269: 18384-18389.
- 9 Kitani A, Nakashima N, Matsuda T, Xu B, Yu S, Nakamura T, and Matsuyama T.
T cells bound by vascular cell adhesion molecule-1/CD106 in synovial fluid in rheumatoid arthritis. Inhibitory role of soluble vascular cell adhesion molecule-1 in T cell activation.
J. Immunol. 1996; 156: 2300-2308.
- 10 Yu S, Nakashima N, Xu BH, Matsuda T, Izumihara A, Sunahara N, Nakamura T, Tsukano M, and Matsuyama T.
Pathological significance of elevated soluble CD14 production in rheumatoid arthritis: in the presence of soluble CD14, lipopolysaccharides at low concentrations activate RA synovial fibroblasts.
Rheumatol. Int. 1998; 17: 237-243.
- 11 Nakashima-Matsushita N, Homma T, Yu S, Matsuda T, Sunahara N, Nakamura T, Tsukano M, Ratman M, and Matsuyama T.
Selective expression of folate receptor β and its possible role in methotrexate transport in synovial macrophages from patients with rheumatoid arthritis.
Arthritis Rheum. 1999; 42: 1609-1616.
- 12 Nakamura T, Baba S, Yamamura Y, Tsuruta T, Matsubara S, Tomoda K, and Tsukano M.
Combined treatment with cyclophosphamide and prednisolone is effective for secondary amyloidosis with SAA1 γ/γ genotype in a patient with rheumatoid arthritis.
Mod. Rheumatol. 2000; 10: 160-164.
- 13 Nakamura T, Yamamura Y, Tsuruta T, Tomoda K, Sakaguchi M, and Tsukano M.
Mycobacterium kansasii arthritis of the foot in a patient with systemic lupus erythematosus.
Intern. Med. 2001; 40: 1045-1049.
- 14 Nakamura T, Tomoda K, Yamamura Y, Tsukano M, Honda I, and Iyama K.
Polyarteritis nodosa limited to calf muscles: a case report and review of the literature.
Clin. Rheumatol. 2003; 22: 149-153.
- 15 Nakamura T, Yamamura Y, Tomoda K, Tsukano M, and Baba S.
Massive hematuria due to bladder amyloidosis in patients with rheumatoid arthritis: three case reports.
Clin. Exp. Rheumatol. 2003; 21: 673-674.
- 16 Nakamura T, Yamamura Y, Tomoda K, Tsukano M, Shono M, and Baba S.
Efficacy of cyclophosphamide combined with prednisolone in a patient with AA amyloidosis secondary to rheumatoid arthritis.
Clin. Rheumatol. 2003; 22: 371-37.
- 17 Nakamura T, Tomoda K, Tsukano M, Yamamura Y, and Baba S.
Gustatory sweating due to autonomic neuropathy in a patient with amyloidosis secondary to rheumatoid arthritis.
Mod. Rheumatol. 2004; 14: 498-501.

- 18 Higashi S, Nakamura T, Tomoda K, Tsukano M, Sakaguchi M, and Kuga F.
A study of ten Japanese patients with seronegative spondylarthropathy: a tentative proposal.
Mod. Rheumatol. 2005; 15: 329-335.
- 19 Nagayoshi R, Nagai T, Matsishita K, Sato K, Sunahara N, Matsuda T, Nakamura T, Komiya S, Onda M, and Matsuyama T.
Effectiveness of anti-folate receptor β antibody conjugated with truncated Pseudomonas exotoxin in the targeting of rheumatoid arthritis synovial macrophages.
Arthritis Rheum. 2005; 52: 2666-2675.
- 20 Nakamura T, Higashi S, Tomoda K, Tsukano M, Baba S, and Shono M.
Significance of SAA1.3 allele genotype in Japanese patients with amyloidosis secondary to rheumatoid arthritis.
Rheumatology (Oxford) 2006; 45: 43-49.
- 21 Migita K, Nakamura T, Maeda Y, Miyashita T, Origuchi T, Yatsuhashi H, Nakamura M, Shibata H, and Eguchi K.
HLA-DRB1*04 alleles in Japanese rheumatoid arthritis patients with AA amyloidosis.
J. Rheumatol. 2006; 33: 2120-2123.
- 22 Nakamura T.
Amyloid A amyloidosis secondary to rheumatoid arthritis: an uncommon yet important complication.
Curr. Rheumatol. Rev. 2007; 3: 231-241.
- 23 Nakamura T, Higashi S, Tomoda K, Tsukano M, and Sugi K.
Primary biliary cirrhosis (PBC)-CREST overlap syndrome with coexistence of Sjogren's syndrome and thyroid dysfunction.
Clin. Rheumatol. 2007; 26: 596-600.
- 24 Horiuchi T, Kiyohara C, Tsukamoto H, Sawada T, Furugo I, Yosizawa S, Ueda A, Tada Y, Nakamura T, Kimoto Y, Mitoma H, Harashima S, Yoshizawa S, Shimada T, Okamura S, Nagasawa K, and Harada M.
A functional M196R polymorphism of tumour necrosis factor receptor type 2 is associated with systemic lupus erythematosus: a case-control study and a meta-analysis.
Ann. Rheuma. Dis. 2007; 66: 320-324.
- 25 Nakamura T, Hirakawa K, Higashi S, Tomoda K, Tsukano M, Iyama K, and Sakae T.
CD8⁺ lymphocytes infiltrate predominantly in the inflammatory foci of MPO-ANCA-positive thoracic hypertrophic pachymeningitis in a patient with HLA-A24.
Mod. Rheumatol. 2007; 17: 75-80.
- 26 Nakamura T, Higashi S, Tomoda K, Tsukano M, and Baba S.
Efficacy of etanercept in patients with AA amyloidosis secondary to rheumatoid arthritis.
Clin. Exp. Rheumatol. 2007; 25: 518-522.
- 27 Nakamura T.
Clinical strategies for amyloid A amyloidosis secondary to rheumatoid arthritis.
Mod. Rheumatol. 2008; 18: 109-118.
- 28 Nakamura T, Higashi S, Tomoda K, Tsukano M, Arizono K, and Nakamura T.
Etanercept treatment in patients with rheumatoid arthritis on dialysis.
Rheumatol. Int. 2010; 30: 1527-1528.
- 29 Migita K, Nakamura T, Koga T, and Eguchi K.
HLA-DRB1 alleles and rheumatoid arthritis-related pulmonary fibrosis.
J. Rheumatol. 2010; 37: 205-207.
- 30 Nakamura T, Higashi S, Tomoda K, Tsukano M, and Shono M.
Etanercept can induce resolution of renal deterioration in patients with amyloid A amyloidosis secondary to rheumatoid arthritis.
Clin. Rheumatol. 2010; 29: 1395-1401.
- 31 Nakamura T.
Amyloid A amyloidosis secondary to rheumatoid arthritis: development of pathophysiology and treatment.
Clin. Exp. Rheumatol. 2011; 29: 850-857.
- 32 Nakamura T.
Amyloid A amyloidosis secondary to rheumatoid arthritis.
In: Amyloidosis-mechanisms and prospects for therapy. (Sarantseva S. eds) Rijeka, Croatia, InTech, 2011; pp23-42.
- 33 Nakamura T, Higashi S, Tomoda K, Tsukano M, and Iyama K.
Cutaneous nodules in patients with rheumatoid arthritis: a case report and review of literatures.
Clin. Rheumatol. 2011; 30: 719-722.
- 34 Migita K and Nakamura T.
TBK1: a potential therapeutic target in RA. New data on a key inflammatory pathway.
Rheumatology (Oxford) 2012; 51: 588-589.
- 35 Migita K, Koga T, Satonuma K, Izumi M, Torigoshi T, Maeda Y, Izumi Y, Jiuchi Y, Mitashita T, Yamasaki S, Aiba Y, Komori A, Nakamura M, Motokawa S, Kawakami A, Nakamura T, and Ishibachi H.
Serum amyloid A triggers the monosodium urate-mediated mature interleukin-1 production from human synovial fibroblasts.
Arthritis Res. Ther. 2012; 14: R119.

- 36 [Nakamura T](#), Higashi S, Tomoda K, Tsukano M, and Shono M.
Effectiveness of etanercept vs cyclophosphamide as treatment of patients with amyloid A amyloidosis secondary to rheumatoid arthritis.
Rheumatology (Oxford) 2012; 51: 2064-2069.
- 37 Furukawa H, Oka S, Shimada K, Sugii S, Ohashi J, Matsui T, Ikenaka T, Nakayama H, Hashimoto A, Takaoka H, Arinuma Y, Okazaki Y, Futami H, Komiya A, Fukui N, [Nakamura T](#), Migita K, Suda A, Nagaoka S, Tsutiya N, and Tohma S.
Association of human leukocyte antigen with interstitial lung disease in rheumatoid arthritis: a protective role for shared epitope.
PLoS One 2012; 7: e33133.
- 38 [Nakamura T](#).
Therapeutic strategies in amyloid A amyloidosis secondary to rheumatoid arthritis.
In: Amyloidosis. (Feng D. eds), Rijeka, Croatia, InTech, 2013; pp213-239.
- 39 [Nakamura T](#), Kumon Y, Hirata S, and Takaoka H.
Abatacept may be effective and safe inpatients with amyloid A amyloidosis secondary to rheumatoid arthritis.
Clin. Exp. Rheumatol. 2014; 32: 501-508.
- 40 [Nakamura T](#), Migita K, Ando Y, Takaoka H, Suzushima H, and Shiraishi N.
Amyloid A amyloidosis in a Japanese patient with familial Mediterranean fever associated with homozygosity for the pyrin variant M694I/M694L.
Mod. Rheumatol. 2014; 24: 349-352.
- 41 [Nakamura T](#) and Migita K.
Prediction of flare-ups in IgG4-related disease. A new concept and call for further elucidation._____
Rheumatology (Oxford) 2015; 54: 5-6.
- 42 Migita K, Izumi Y, Agematsu K, Matsumoto J, Jiuchi Y, Kozuru H, Nonaka F, Shimizu T, [Nakamura T](#), Iwanaga N, Furukawa H, Yasunari M, Kawakami A, and Eguchi K.
Dysregulated mature IL- β production in familial Mediterranean fever.
Rheumatology (Oxford) 2015; 54: 660-665.
- 43 [Nakamura T](#), Ueda M, Takaoka H, and Ando Y.
Trigeminal sensory neuropathy and mixed connective tissue disease: a case report.
Clin. Case Rep. Rev. 2015; 1: 130-132.
- 44 [Nakamura T](#).
Mycobacterium chelonae infection and TNF α blockade.
Med. Sci. Rev. 2015; 2: 15-18.
- 45 Oka S, Furukawa H, Shimada K, Sugi S, Hashimoto A, Komiya A, Fukui N, Suda A, Tsunoda S, Ito S, Katayama M, [Nakamura T](#), Saisho T, Sano H, Migita K, Nagaoka S, Tsutiya N, and Tohma S.
Association of human leukocyte antigen alleles with chronic lung diseases in rheumatoid arthritis.
Rheumatology (Oxford) 2016; 55: 1301-1307.
- 46 [Nakamura T](#).
Developments in the treatment of amyloid A amyloidosis secondary to rheumatoid arthritis.
In: Exploring new findings on amyloidosis. (Fernandes-Escamilla A-M eds), Rijeka, Croatia, InTech, 2016; pp215-241.
- 47 [Nakamura T](#), Hirakawa K, Takaoka H, and Iyama K.
Dystrophic calcinosis with both a huge calcified mass in the cervical spine and calcification in the chest wall in a patient with rheumatoid overlap syndrome.
Clin. Rheumatol. 2016; 35: 1403-1409.
- 48 [Nakamura T](#) and Yamamoto T.
Potential of a 70kDa IL-10-like factor in synovial fluid from rheumatoid arthritis patients to augment superoxide generation by human neutrophils.
Int. J. Rheum. Dis. 2016; 19: 150-158.
- 49 Yahiro M, Furukawa H, Asano T, Sato S, Kobayashi H, Watanabe H, Suzuki E, [Nakamura T](#), Koga T, Umeda M, Nonaka F, Ueki Y, Eguchi K, Kawakami A, and Migita K.
Serum amyloid A1 (SAA1) gene polymorphism in Japanese patients with adult-onset Still's disease.
Medicine (Baltimore) 2018; 97: 49 (e13394).
- 50 Otsubo H, Tsuneyoshi Y, [Nakamura T](#), Matsuda T, Komiya S, and Matsuyama T.
Serum-soluble folate receptor b as a biomarker for the activity of rheumatoid arthritis synovitis and the response to anti-TNF agents.
Clin. Rheumatol. 2018; 37: 2939-2945.
- 51 [Nakamura T](#).
Protein-losing enteropathy may be an important characteristic manifestation in Sjogren's syndrome.
Mod. Rheumatol. 2019; 29: 397-399.
- 52 [Nakamura T](#), Shiraishi N, Morikami Y, Fujii H, and Kuratsu J.
Amyloid A amyloidosis secondary to rheumatoid arthritis may be treatable but is still difficult to manage in daily clinical practice.
Amyloid 2019; 26 (Supl 1): 123-124.

- 53 [Nakamura T](#), Shiraishi N, Morikami Y, Fujii H, Itoshima H, and Kamio T.
Protein-losing enteropathy may be an important characteristic manifestation in Sjogren's syndrome.
Mod. Rheumatol. 2019; 29: 397-399.
- 54 Taylor PC, Saurigny D, Vencovsky J, Takeuchi T, [Nakamura T](#), Matsievskaja G, Hunt B, Wanger T, and Souberbielle; NEXUS Study Group.
Efficacy and safety of namilumab, a human monoclonal antibody against granulocyte-macrophage colony stimulating factor (GM-CSF) ligand in patients with rheumatoid arthritis (RA) with either an inadequate response to background methotrexate therapy or an inadequate response or intolerance to an anti-TNF (tumour necrosis factor) biologic therapy: a randomized, controlled trial.
Arthritis Res. Ther. 2019; 21: 101 1186/s13075-019-1879-x.
- 55 [Nakamura T](#), Shiraishi N, Morikami Y, Fujii H, and Yoshinaga T.
Amyloid A amyloidosis in a patient with Caplan's syndrome, with special reference to genetic predisposition.
Mod. Rheumatol. Case Rep. 2020; 4: 212-217.
- 56 [Nakamura T](#).
Coexistence of anti-SRP and anti-SS-A/Ro antibodies in inflammatory myopathy: does the association occur by chance? A case report.
S.N. Comprehensive Clin. Med. 2020; 2: 822-828.
- 57 Imamura M, Mukaino A, Takamuatsu K, Tuboi H, Higuchi O, Nakamura H, Abe S, Ando Y, Matsuo H, [Nakamura T](#), Sumida T, Kawakami A, and Nakane S.
Ganglionic acetylcholine receptor antibodies and autonomic dysfunction in autoimmune rheumatic diseases.
Int. J. Mol. Sci. 2020;21: 1332-1347.
- 58 Goto T, Miyazaki Y, Nakayamada S, Shiraishi N, Yoshinaga T, Tanaka Y, and [Nakamura T](#).
Down-regulated Th17 cells in a patient with familial Mediterranean fever associated with amyloid A amyloidosis in the treatment of canakinumab.
Mod. Rheumatol. Case Rep. 2023; 7: 237-242.
- 59 Higuchi T, Oka S, Furukawa H, Sumida K, Tsunoda S, Ito S, Okamoto A, Fujimori M, [Nakamura T](#), Katayama M, Saisho K, Shinohara S, Matsui T, Migita K, Nagaoka S, and Tohma S.
Association of a single nucleotide variant in TERT with airway disease in Japanese rheumatoid arthritis patients.
Genes. 2023; 14: 2084-2092.
- 60 Higuchi T, Oka S, Shimada K, Tsunoda S, Ito S, Okamoto A, Fujimori M, [Nakamura T](#), Katayama M, Saisho K, Shinohara S, Matsui T, Migita K, Nagaoka S, Tohma S, and Furukawa H.
Association of rare single nucleotide variant MUC5B rs356705950 with interstitial lung disease in Japanese rheumatoid arthritis.
Rheumatology (Oxford) 2024; Aug 7: keae422. doi: 10.1093/rheumatology/keae422.
- 61 Matsuki Y and [Nakamura T](#).
Spondyloarthritis and Tietze's syndrome: a re-evaluation.
Mod. Rheumatol. 2025; 35: 1-6.